

دو گونه و یک گزارش جدید از سرده گون (تیره باقلانیان) از شمال شرق ایران

یاسمین ناصح* و محمدرضا جوهرچی

دریافت: ۱۳۹۶/۰۷/۲۲ / ویرایش: ۱۳۹۷/۰۴/۱۴ / پذیرش: ۱۳۹۷/۰۴/۲۴ / انتشار: ۱۳۹۷/۰۹/۳۰

گروه گیاه شناسی، پژوهشکده علوم گیاهی، دانشگاه فردوسی مشهد، مشهد، ایران

*مسئول مکاتبات: Naseh-ya@staff.um.ac.ir

چکیده. ضمن بررسی نمونه‌های گیاهی جمع آوری شده متعلق به بخش *Ammodendron* از سرده *Astragalus* موجود در هرباریوم پژوهشکده علوم گیاهی دانشگاه فردوسی مشهد دو گونه جدید به اسامی *Astragalus rashed-mohasseli* و *Astragalus microfoliolatus* از این منطقه کشف و برای نخستین بار برای دنیا پیشنهاد و معرفی می‌شوند. شواهد ریخت‌شناسی تعلق این گونه‌ها را به این بخش نشان می‌دهد. گونه *Astragalus microfoliolatus* به دلیل کوچک بودن ابعاد برگچه‌ها که از ویژگی‌های شاخص آن به شمار می‌رود از گونه‌های خویشاوند خود جدا می‌شود. از طرف دیگر گونه *Astragalus rashed-mohasseli* با حضور کرک روی بخش پشتی درفش گل که برای نخستین بار در یکی از گونه‌های این بخش مشاهده می‌شود، مادگی مستطیلی کشیده و محور برگگی کوتاه از گونه خویشاوند خود متمایز می‌شود. این گونه‌ها از شمال شرقی ایران، استان‌های خراسان رضوی و جنوبی جمع‌آوری شده‌است و در هرباریوم پژوهشکده علوم گیاهی دانشگاه فردوسی مشهد نگهداری می‌شود. همچنین گونه *A. aiwadzhi* گزارش جدیدی از این منطقه برای فلور ایران است. شرح گونه‌های جدید، نقشه انتشار و همچنین قرابت آنها با نزدیکترین خویشاوندانشان به همراه تصاویر گونه‌ها در این مقاله ارائه گردیده‌است.

واژه‌های کلیدی. آمودندرون، استان‌های خراسان، ایرانوتورانی، تاکسونومی، صفات ریختی

Two new species and a new record of the genus *Astragalus* (Fabaceae) from NE Iran

Yasamin Nasseh* & Mohammad Reza Joharchi

Received: 14.10.2017 / Revised: 05.07.2018 / Accepted: 15.07.2018 / Published: 21.12.2018

Research Center for Plant Sciences, Ferdowsi University of Mashhad, Mashhad, Iran

*Correspondent author: Naseh-ya@staff.um.ac.ir

Abstract. During the study on the specimens of *Astragalus* sect. *Ammodendron* in the herbarium of Ferdowsi University of Mashhad (FUMH), based on the last and most recent sources, two new species, i.e. *A. microfoliolatus* sp. nova and *A. rashed-mohasseli* sp. nova were found and described. Morphological evidence confirmed that the belonging of these taxa to *Astragalus* sect. *Ammodendron*. The most important differences between *A. microfoliolatus* and its closest relatives are leaflets with very small size and glabrescent legume. Also, the main distinctive characters of *A. rashed-mohasseli* are the presence of hairs on the dorsal side of vexillum, oblong legume and short rachis. Hairy vexillum has been observed for the first time in this section. In addition, *A. aiwadzhi* is recorded as a new species for the flora of Iran. Taxonomic descriptions, illustrations and distribution maps are provided to expedite identification.

Keywords. *Ammodendron*, Irano-Turanian, Khorassan provinces, morphological characters, taxonomy

INTRODUCTION

Astragalus L. (Linnaeus, 1753) is the largest genus of flowering plants with approximately 3000 species worldwide. Section *Ammოდendron* Bunge (Bunge, 1868, 1869) is one of the most complex groups among those covered with bifurcate hairs. *Astragalus* sect. *Ammოდendron* was first established by Bunge with 19 species. According to recent studies, however, this group includes approximately 80 species worldwide of which 35-40 species and 25 endemics are distributed in Iran (Rechinger *et al.*, 1961; Maassoumi, 1998, 2005; Podlech *et al.*, 2010; Ghahremaninejad, 2004a,b, 2005; Ghahremaninejad & Gaskin, 2004; Nasseh *et al.*, 2010, 2012; Dastpak *et al.*, 2011; Podlech & Zarre, 2003, 2013). Iran is one of the most genetically diverse hotspots of *Astragalus* with about 826 species (Ghahremaninejad, 2015).

The plants in this section are shrubby or sub-shrubby, often becoming more or less leaflet-less with age. After the development of the stems, they are densely covered with appressed medifixed, white hairs. Stipules are often membranaceous, somewhat adnate to the petiole or almost free. Petioles and rachises are sometimes hard and spiny. Leaflets are found in 1-3 pairs, rarely unifoliate. Inflorescences are usually lax with short or long peduncles. Bracteoles are absent. Calyces are campanulate or nearly so and not inflated in fruit. Pods coriaceous, include 2 locules, slightly longer than the calyx, sessile to stipitate, valves densely covered with soft, long, appressed or spreading, basifixed hairs, sometimes mixed with short medifixed hairs or rarely glabrous (Podlech & Zarre, 2013).

Phytogeography. The diversity center of *Astragalus* section *Ammოდendron* is the Turkestanian floristic Province (Takhtajan, 1986) of the Irano-Turanian region. The members of this group are vastly distributed in the Armeno-Iranian Province of the aforementioned region. The great majority of species of sect. *Ammოდendron* (Maassoumi, 1998; Podlech & Zarre, 2003, 2013; Podlech *et al.*, 2010) occurs in Middle Asia, Egypt (Sinai), Turkey, Saudi Arabia, Near East, Caucasus region, Iran (mainly in the central, southern and eastern parts of Iran) and Afghanistan and Pakistan (Ghahremaninejad, 2004b; Podlech & Zarre, 2013). They are mostly psammophytes or associated with more or less sandy substrata, in dry and semidesertic regions and very rarely petrophilous plants of foothills and low mountains. (Ghahremaninejad, 2004b; Podlech & Zarre, 2013).

The aim of the present work is to study the taxonomy of the genus *Astragalus* section *Ammოდendron* with a focus on collected materials from NE Iran. Since Middle Asia in Irano-Turanian region is the main diversity center of *Astragalus* sect. *Ammოდendron* (Komarov, 1946; Maassoumi, 1998), the vicinity of Razavi Khorassan province to this area resulted in the occurrence of many species of this section in NE Iran.

MATERIALS AND METHODS

Herbarium specimens of the genus *Astragalus* sect. *Ammოდendron* in the herbarium of Ferdowsi University of Mashhad, (FUMH) were revised. Studies were carried out by using different floras and resources (Gontscharov, 1946; Maassoumi, 2005; Podlech & Zarre, 2010, 2012, 2013) and related taxonomic literatures (Bunge, 1868, 1869; Gontscharov, 1946; Rechinger *et al.*, 1961; Maassoumi, 1998, 2005; Podlech *et al.*, 2010; Ghahremaninejad, 2004a,b, 2005; Ghahremaninejad & Gaskin, 2004; Nasseh *et al.*, 2010, 2012; Dastpak *et al.*, 2011; Podlech & Zarre, 2003, 2013). The most important characteristics of the examined specimens were the size and shape of leaflets, the characteristics of inflorescence comprising the length of peduncle and raceme, the size of calyces and the type and color of their hairs, the number of flowers, the shape and size of the legume and their indumentum types.

RESULTS AND DISCUSSION

Taxonomic treatment

New taxa to the world

1. *Astragalus microfoliolatus* Nasseh, *sp. nova* (Figs. 1, 2; Tables 1, 2)

Type: Iran. Razavi Kohrassan: N Torbat-e Heydariyeh, Robatsang, 4 October 1991, 1650 m, Joharchi & Zangooei 21038! (Holotype: FUMH!; Isotype: T!).

Diagnosis: Differt ab *A. squarrosus* Bunge: *Foliis* 2-5 × 1.5-2 mm (*nec* 5-12 × 1.5-4 mm). *Rachis* 0.2-0.7 cm *longi* (1-4 cm). *Bracteae* 0.5 mm *longi* (*nec* 1-1.5 mm). *Calyx* 5-6 mm *longi*, *subtus pilis obsita paulatim glabrescens* (*nec* 6-8 mm, *dense pilosa*). *Leguminibus* 4.5-5.5 mm *longi*, *subtus pilis obsita paulatim glabrescens* (*nec* 6-8 mm, *dense pilosa*).



















Fig. 1. *A. microfoliolatus* Nasseh; Joharchi & Zangoeie 21038 (Photo by H. R. Sharghi).

Table 1. Comparison of *A. microfoliolatus* & *A. squarrosus*

species character	<i>A. microfoliolatus</i>	<i>A. squarrosus</i>
Leaflet	2-5 × 1.5-2	5-12 × 1.5-3(-4)
Rachis (cm)	0.2-0.7	1-3(-4)
Bract (mm)	0.5	1-1.5
Calyx (mm)	5-6	6-8
	covered with sparse hairs gradually glabrescent	covered with dense hairs

Table 2. Some floral and vegetative parts of the new *Astragalus* species

Leaflet	legume	Ovary	Stamen	Carina	Wing	Standard	Calyx	Species
								<i>A. microfoliolatus</i>
								<i>A. rashed-mohasseli</i>

Description: Plants perennial, up to 35 cm tall, suffruticose, branched, densely covered with medifixed appressed white hairs. Stipules 1-1.5 mm long, membranous, shortly adnate to the petiole, sparsely appressed white hairy. Rachis very short up to 7 mm long; leaflets mainly 2 pairs rarely 1, oblong to elliptic, 2-4 × 1.5-2 mm, mainly acute, rarely obtuse, on upper side subglabrous to glabrous, on underside densely hairy. Peduncles 0.5-1.5 cm long, densely covered with medifixed appressed hairs. Racemes 3-13 cm long, covered with medifixed appressed hairs, 4-12 flowered. Bracts ca. 0.5 mm long, membranous, ovate, subglabrous. Pedicel ca. 1.5 mm long, white hairy. Calyx 5-6 mm long, campanulate to tubular, covered with medifixed appressed white hairs; teeth short triangular, 1 mm

long, hairy on outer and inner side. Corolla yellow when drying; standard ca. 11 mm long, elliptic, wings 10.5-11 mm long, limb ca. 5 mm long, oblong, claw 5.5-6 mm long, auricle small, less than 0.5 mm long; keel 10-1.5 mm long; limb ca. 5 mm long, elliptic-oblong, claw ca. 5 mm long, auricle very short. Legumes ca. 5 mm long, ovate, covered with basifixed spreading hairs on little tubercles, without short hairs, with a short glabrous stipe ca. 1 mm long, glabrous and a short beak ca. 0.75 mm long. Flowering and Fruiting period: May-June.

Paratype: Iran. Razavi Khorassan: Torbat-e Heydariyeh-Mashhad road, Robatsang, 20.8.1986, Ayatollahi & Zangooei 14998! (FUMH).

Discussion: *Astragalus microfoliolatus* is a new endemic species, its distribution being restricted to a limited mountainous area with a temperate to arid climate in Razavi Khorassan, NE Iran. It has been collected only from the type locality. This species is very close to *A. squarrosus*, with some differences in morphological characters summarized in Table 3. On the basis of IUCN Red List categories and criteria

(IUCN, 2010) this species is classified as critically endangered. Therefore, a conservation program for this species is highly recommended.

Etymology: The name of the *Astragalus microfoliolatus* for the new species is proposed due to its very little leaflets.

Table 3. Comparison of *A. rashed-mohasseli* & *A. macrobotrys*

species character	<i>A. rashed-mohasseli</i>	<i>A. macrobotrys</i>
Ovary (mm)	8-9 × 1-1.5, oblong, covered with tangled homomorphic hairs	5-7 × 2, ellipsoid, covered with heteromorphic, long and short hairs
Rachis (cm)	0.2-1	1.5-3
Standard	hairy, limb rumboid	glabrous, limb obovate and rotundate at the apex



Fig. 2. Distribution Map: ● *Astragalus microfoliolatus*, ▲ *Astragalus rashed-mohasseli*, ■ *Astragalus aiwadzhi*.

2. *Astragalus rashed-mohasseli* Nasseh, sp. nova (Figs. 2, 3; Tables 2, 3)

Type: **Iran.** Razavi Khorassan, SE Kalat-e Naderi, between Hamam-Ghale & Baba-Faraji, N 36° 52' 28.31 " E 59° 53' 54.26", 970 m, 10 June 2009, Joharchi 42985 (Holotype: FUMH!; Isotype: T!).

Diagnosis: Differt ab *A. macrobotrys* Bunge: *Rachis* 0.2-1 cm longi (nec 1-3.5 cm). *Vexillum* rhomboid, *pilosi* (nec obovato, apice rotundatum, non glabrescens). *Ovary* 8-9 mm oblonga (nec 5-7 mm, ellipsoidea).



Fig. 3. *A. rashed-mohasseli* Nasseh; Joharchi 42985 (Photo by H.R. Sharghi).



Fig. 4. *A. aiwadzhi* B.Fedtsch.; Ayatollahi 15063 (Photo by H.R. Sharghi).

Description: Plants perennial, up to 65 cm tall, suffruticose, branched, covered with medifixed appressed white hairs. Stipules 1.5-4 mm long, membranous, adnate to the petiole, somewhat vaginate and connate, nearly with free tips triangular, sparsely hairy or subglabrous. Rachis 2-10 mm long. Leaflets in 1-2 pairs, 7-15 × 3-7 mm, elliptic to obovate, rounded at apex, on upper side sparsely hairy to subglabrous, on underside densely medifixed appressed hairy. Peduncles (1-) 3-7 cm long. Racemes (3-) 5-20 cm long when flowering, (8-) 10-30 flowered. Bracts 1-2 mm long, ovate, membranous, very sparsely white and black hairy, sometimes sparsely ciliate at the margins. Pedicels 1-2 mm long, white hairy. Calyx 4.5-5.5 mm long, campanulate-turbinate, covered with medifixed, appressed white hairs, very rarely mixed sparsely black hairs; teeth 1-2 mm long, narrowly triangular to triangular, sparsely hairy on inner side. Corolla yellow to pale violet when drying. Standard ca. 12 mm long, limb 6.5-7 mm long, rhombic-elliptic, plicate at margin, hairy on outer side in upper part, claw 5 mm long; wings 10.5-11 mm long, limb ca. 5 mm long, oblong, claw 6-6.5 mm long, auricle up to 0.5 mm; keel 10 mm long limb ca. 5 mm long, claw 5.5 mm. Stamen 9.5-10 mm long. Ovary oblong, substipitate, only covered with long, tangled hairs. Legume unknown but based on the ovary character, it might be oblong.

Flowering and Fruiting period: May-June

Discussion: *Astragalus rashed-mohasseli* is a new species which has been collected from Kalat-e Naderi, a small part of Razavi Khorassan, NE Iran. This type locality is a temperate mountainous area. Collected only from the type locality mentioned before, this species is very close to *A. macrobotrys*, being different from the latter in the characteristic of their standards due to the existence of hairs on the former (Table 1). On the basis of IUCN Red List categories and criteria (IUCN 2010) this species is classified as critically endangered, therefore, a conservation program for this species is highly recommended.

Etymology: This species is dedicated to Professor Dr. Mohammad-Hassan Rashed-Mohassel (Ferdowsi University of Mashhad), the former Head of the FUMH herbarium.

New Record to Iran

Astragalus aiwadhzi B.Fedtsch., Not. Syst. Leningrad 8: 166 (1940). (Figs. 2, 4)

Iran. Razavi Khorassan: 21 km of Sarakhs towards Sangar, N 36° 21' 7.1994"E 61° 8' 13.2"30 March 1987, Ayatollahi 15063! (FUMH).

Description: Plants perennial, up to 55-65 cm tall, subshrub, branched; ligneous older branches up to 25 cm long, grayish; branches of the current year 20-55 cm long. Stipules 2-2.5 mm long, membranous, totally connate, bidentate sparsely hairy or subglabrous. Rachis 1.5-3 cm long. Leaflets in lower parts usually in two pairs, in upper parts in one pair, 1.5-4.5 cm × 2-3 mm, linear acute densely covered on both sides with appressed white hairs. Peduncles 1.5-3 cm long. Racemes loose, 10-20 cm long with 15-20 flowers. Bracts lanceolate 1.5 (-2) mm long, ovate, membranous, very sparsely white ciliate. Pedicels 1-1.5 mm long, white hairy. Calyx 6-7 mm long, campanulate-cylindric, covered with medifixed, appressed white hairs, very rarely mixed sparsely black hairs; teeth 1-1.5 mm long, narrowly triangular to triangular, sparsely hairy on inner side. Corolla yellow to pale violet when drying. Standard 11-12 mm long, limb rounded-oval, side in upper part claw 5 mm long; wings 10-11 mm, limb oblong; keel 9-10 mm long. Legumes narrowly ellipsoid, 6-8 mm long, 2-3 mm high, with a short beak and a stipe 0.5-1 mm long; valves covered with spreading white long and short hairs.

This species has been described and introduced from Tadjikistan in ditone Schaartus, Vallis Bischkent, inter Pagos Aiwadzheth Czaschmy (Gontscharov, 1946). Here we report it as a new record for the first time from Iran.

ACKNOWLEDGEMENT

We would like to appreciate Research Council of Ferdowsi University of Mashhad for their financial support of the research project no. 851. Also, we sincerely thank Hamid-Reza Sharghi for the photographs.

REFERENCES

- Bunge, A. 1868-1869. Generis Astragali species Gerontogae. Pars prior, Claves Diagnosticae. Mémoires de l'Académie Impériale des sciences de Saint Petersburg. Septième série. 7, 11: 1-140. Pars altera, Specierum enumeratio., ser. 11 (16): 129 (1868); I. C. 15 (1): 2269 p.
- Dastpak, A., Maassoumi, A.A., Kazempour Osaloo, S., and Nejadstari, T. 2011. *Astragalus ghamishluensis* (Fabaceae), a new species of sect. *Ammodendron* from Iran. – Iran J. Bot. 17: 175-177.
- Ghahremaninejad, F. 2004a. Value of trichom characters for separation of bifurcating hairy *Astragalus* L. (Fabaceae) at the sectional level. – Turk. J. Bot. 28: 241-245.
- Ghahremaninejad, F. 2004b. The sections of *Astragalus* L. with bifurcating hairs in Iran. – Turk. J. Bot. 28: 101-117.

- Ghahremaninejad, F.** 2004c. *Astragalus baharensis* (Fabaceae), a new species from NE Iran. – Ann. Bot. Fenn. 41: 143-145.
- Ghahremaninejad, F. and Gaskin, J.F.** 2004. A new species of *Astragalus* (Fabaceae, Atari be Gakegeae) from Iran. – Novon. 14: 431-433.
- Ghahremaninejad, F.** 2005. *Astragalus hekmat-safaviae* (Fabaceae), a new species from Iran. – Ann. Bot. Fenn. 42: 313-315.
- Ghahremaninejad, F.** 2015. Notes about *Astragalus* (Leguminosae) in Iran. – Ann. Naturhist. Mus. Wien. B 117: 279-281.
- Gontscharov, N.F.** 1946. *Astragalus*. – In Komarov, V.L. (ed.) Flora of the U.S.S.R. vol. 12. – Moskova and Leningrad. 681 p.
- Komarova, V.L.** 1940. Botanicheskie materialy gerbariya botanicheskogo Instituta Imeni Akademii Nauk S S R. Leningrad. Leningrad [St. Petersburg] vol. 8: 166.
- Linnaeus, C. von.** 1753. Species Plantarum, vol. 2. Laurentius Salvius, Stockholm. 755 p.
- Maassoumi, A.A.** 2005. The genus *Astragalus* in Iran, vol. 5 Research Institute of Forests and Rangelands, Tehran 786 p.
- Maassoumi A.A.** 1998. Old world check-list of *Astragalus*. Research Institute of Forests and Rangelands, Tehran 617 p.
- Nasseh, Y., Joharchi, M.R. and Maassoumi, A.A.** 2010. Two new species of the genus *Astragalus* sect. *Ammოდendron* (Fabaceae) from Iran. – Iran J. Bot. 16: 221-224.
- Nasseh, Y. and Joharchi, M. R.** 2012. New records and new synonym of *Astragalus* sections *Ammოდendron* and *Caprini* (Fabaceae) from Iran. – Iran J. Bot. 18 (1): 76-83.
- Podlech, D. and Zarre, Sh.** 2003. New species of *Astragalus* sect. *Ammოდendron* (Fabaceae) with a key for the species of the Flora Iranica area. – Willdenowia. 33: 314-352.
- Podlech, D., Zarre, Sh., Maassoumi, A.A., Ekici, M. and Sytin A.** 2010. *Astragalus* sect. *Ammოდendron*. – In Rechinger, K.H. (ed.), Flora Iranica. vol. 178: 373-403. – Naturhistorischen Museum, Wien.
- Podlech, D., Maassoumi, A.A. and Zarre, Sh.** 2012. *Astragalus* sect. *Ammოდendron*. – In Rechinger, K.H. (ed.), Flora Iranica. vol. 179, 312 p. – Naturhistorischen Museum Wien.
- Podlech, D. and Zarre, Sh.** 2013. A taxonomic revision of the genus *Astragalus* L. (Leguminosae) in the Old World. vols. 1-3. Naturhistorisches Museum Wien. 2439 p.
- Rechinger, K.H., Dulfer, H. and Patzak, A.** 1961. Sirjaevii fragmenta astragalologica XV. *Astragalus* sect. *Ammოდendron*-Sitzungsber. Oesterr. Akad. Wiss., Mathem. – Natur. Kl., Abt. 170: 35-53.

How to cite this article:

Nasseh, Y. and Joharchi, M. R. 2018. Two new species and a new record of the genus *Astragalus* (Fabaceae) from NE Iran. – Nova Biol. Reperta 5 (3): 311-319.