

## دو گونه جدید از سرده کوزینیا (تیره کاسنیان) از غرب و شمال شرق ایران

فریده عطار<sup>۱</sup>، حسین معروفی<sup>۲</sup>، منصور میرتاج‌الدینی<sup>۳</sup> و آرش ستوده<sup>۴</sup>

<sup>۱</sup> هرباریوم مرکزی دانشگاه تهران، دانشکده زیست‌شناسی، دانشکده‌گان علوم، دانشگاه تهران، ایران؛ <sup>۲</sup> بخش تحقیقات جنگل‌ها و مراتع، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی کردستان، سازمان تحقیقات، آموزش و ترویج کشاورزی، سنندج، ایران؛ <sup>۳</sup> گروه زیست‌شناسی، دانشکده علوم، دانشگاه شهید باهنر کرمان، کرمان، ایران؛ <sup>۴</sup> بخش تحقیقات آگرواکولوژی، دانشگاه برگندی، دیژون، فرانسه.  
مسئول مکاتبات: آرش ستوده arash.sotoodeh@u-bourgogne.fr

چکیده. دو گونه جدید از سرده کوزینیا شرح داده می‌شود. کوزینیا برنمولریانا از غرب ایران که بر اساس بعضی صفات کلیدی مانند تعداد گل و برگک و شکل گل‌آذین در بخش‌هاوسکنختی قرار می‌گیرد. این گونه با گونه نزدیکش یعنی کوزینیا هاوسکنختی و کوزینیا کرکسنسیس مقایسه می‌شود. دومین گونه جدید به نام کوزینیا سمنانسیس از شمال شرق ایران شرح داده می‌شود که بر اساس صفاتی چون تعداد گل، ساقه بالدار و شکل برگک در بخش‌هاستونسفاله قرار می‌گیرد و با نزدیکترین خویشاوندان خود یعنی گونه‌های کوزینیا الکسینکوانا و کوزینیا گلوکوپسیس مورد مقایسه قرار می‌گیرد. عکس‌های این گونه‌ها و نقشه‌های پراکنش‌ارایه می‌شوند.

واژه‌های کلیدی. کوزینیا، آرایه‌شناسی، ریخت‌شناسی، بورنمولر، فلور ایران

## Two new species of genus *Cousinia* (Asteraceae, Cardueae) from W and NE Iran

Farideh Attar<sup>1</sup>, Hossein Maroofi<sup>2</sup>, Mansour Mirtadzadini<sup>3</sup> & Arash Sotoodeh<sup>4</sup>

<sup>1</sup> Central herbarium of Tehran University, School of Biology, College of Science, University of Tehran, Tehran, Iran; <sup>2</sup> Forests and Rangelands Research Department, Kurdistan Agricultural and Natural Resources Research and Education Center, AREEO (Agricultural Research, Education & Extension Organization), Sanandaj, Iran; <sup>3</sup> Department of Biology, Faculty of Science, Shahid Bahonar University of Kerman, Kerman, Iran; <sup>4</sup> Agroécologie, INRAE, Institut Agro, Univ. Bourgogne, Univ. Bourgogne Franche-Comté, F-21000 Dijon, France.  
Correspondent author: Arash Sotoodeh, arash.sotoodeh@u-bourgogne.fr

**Abstract.** Two new species of *Cousinia* are described from Iran. *Cousinia bornmulleriana* from the west that can be placed in sect. *Haussknechtii* based on some key characters such as flower numbers, phyllary numbers and the inflorescence form traits. This species can be also compared with its closest relative *C. haussknechtii*. and *C. karkasensis*. The second new species, *C. semnanensis*, from North East of Iran, belongs to the sect. *Stenocephalae* for its flower numbers, winged stem, and phyllaries shape, and is compared to *C. alexeenkoana* and *C. glaucopsis* as its close relative. Natural photos and distribution maps are presented.

**Keywords.** *Cousinia*, taxonomy, morphology, Bornmüller, flora of Iran

## INTRODUCTION

The genus *Cousinia* Cassini is one of the largest genera of flowering plants, with 600–700 species distributed in central and western Asia (López-Vinyallonga et al., 2009; Rechinger, 1972, 1979, 1986; Sennikov, 2013; Tscherneva, 1962; Atazadeh et al., 2020).

In Iran the Kopet Dag, Alborz Mts., Zagros Mts., and NW regions are the main distribution centers of this genus. Most species have a limited distribution range and occupy small isolated habitats (Attar & Djavadi, 2010). In the desertic area of Central Iran, members of *Cousinia* are completely absent except for some such as *C. piptocephala* Bunge and *C. deserti* Bunge.

Despite the Irano-Turanian nature of most taxa, a few ones are presented in the transition areas between Hyrcanian and Irano-Turanian phytogeographical regions. Some of these species include *C. pterocaulos* (C.A.Mey.) Rech.f., (Sect. *Serratuloides* Bunge), *C. chamaepeuce* Boiss. (Sect. *Sphaerocephalae* Bunge). A few species grow near the Sahara–Sindian and Sudanian regions, some of them are *C. stocksii* C.Winkl. (Sect. *Stocksianae* Rech.f.) and *C. gedrosiaca* Bornm. & Gauba (sect. *Lachnosphaerae* Rech.f.).

Based on Flora Iranica (Rechinger, 1972, 1979), there are about 220 taxa for the flora of Iran. In recent years many species described or recorded for the first time for the flora of Iran (Tscherneva et al., 2005; Djavadi & Attar, 2006; Attar et al., 2007, 2016; Assadi, 2009, 2010, 2011; Attar & Mirtadzadini, 2009; Attar & Djavadi, 2010; Mehregan & Assadi, 2010; Mehregan et al., 2010; Attar, 2011; Mirtadzadini & Attar, 2014; Rastegar et al., 2018; Attar & Amini Rad, 2019).

Also, according to Flora Iranica (Rechinger, 1972, 1979), *Cousinia* is divided into 71 sections of which 42 sections are present in Iran. Recently, three sections including *Ramosissimae* Djavadi & Attar (Djavadi & Attar, 2006), *Argentae* Mehregan & Assadi (Mehregan & Assadi, 2010), and *Chaetocephalae* Assadi (Assadi, 2010) have been described or validated.

Some useful characters for delimiting different sections of *Cousinia* taxa are including being annual or perennial, glabrous or hairy anthers, ± decurrent stem leaves, leaves segments, corolla color, number of flowers in each capitulum, number of flowers, synflorescence, appendiculate or exappendiculate phyllaries, and a smooth or spinulose margin of phyllaries.

Among collected *Cousinia* materials from Kordestan and Semnan provinces, we found two interesting specimens that were different from the other taxa by some distinct characters. Therefore, we

visited the same localities again in the year after, and more materials were collected. Among the examined herbarium specimens, we also noticed that the specimen from Herbarium Haussknecht (JE), accompanied by an identification note by Bornmüller, has been identified as a form of *C. haussknechtii* f. *ramosa* in one herbarium sheet, and *C. haussknechtii* f. *subspicata* in the other sheet (Fig. 1). After studying and comparing our specimens with different herbarium specimens and related literature, they are recognized as new species.

## MATERIAL AND METHODS

During the study of the genus *Cousinia* and several travels by the authors to the different regions of Iran, two new *Cousinia* species were found. The collected specimens were accurately examined and compared with their related species in different herbaria mostly in TUH, TARI, MIR, G, and JE. In addition, all the type specimens were compared with these two new species. We also used Flora Iranica (Rechinger, 1972, 1979), Flora of Turkey (Huber-Morath, 1975), Flora of the U.S.S.R. (Tscherneva, 1962), and recent related publications as literature sources.

## TAXONOMY

### Sect. *Haussknechtianae* Rech.f.

Updated and completed description.

Perennial, tall, cauline leaves leathery, not decurrent, almost amplexicaul, discolored, or concolored. Synflorescence a large panicle. Capitula ca. 20–60-flowered; phyllaries 60–160, exappendiculate, acicular, imbricate or recurved; bristles smooth.

### *Cousinia bornmulleriana*

Attar & Maroofi, **sp. nov.** (Fig. 2)

**Diagnosis.** — *Cousinia bornmulleriana* is related to sect. *Haussknechtianae* by its capitulum shape and discolored lanceolate leathery leaves. (Table 1). We also noticed in the specimen of JE herbarium that Bornmüller recognized the differences between this specimen and *C. haussknechtii*, so he identified this specimen as *C. haussknechtii* f. *ramosa* (Fig. 1).

**TYPE:** Iran, Kordestan Province: 30 km from Sanandaj to Kamyaran, Dulab pass, Avallan mountain, 1950 m, 27.7.2009, Attar, Maroofi & Zamani FA60001 (Holotype, TARI; Isotype, IRAN).

Perennial, rootstock woody. Stem more or less 80 cm long, erect, ivory white, at first arachnoid, then gradually glabrescent, leafy in the lower half, divaricately branched in the upper half, forming a long corymbose synflorescence. Leaves rigid, leathery,



**Figure 1.** Left and middle: *Cousinia haussknechtii* f. *ramosa* and Right: *C. haussknechtii* f. *subspicata*; identified as different forma by Bornmüller (images from herbarium Haussknecht, JE).



**Figure 2.** Left: *Cousinia bornmulleriana* Attar & Maroofi, sp. nov. (Holotype, Attar, Maroofi & Zamani FA60001); Right: Holotype specimen of *C. haussknechtii*, Haussknecht 569, from Genève Herbarium.

discolored, green, loosely arachnoid on the upper surface, white densely arachnoid beneath, venation pinnate-reticulate, distinctly prominent on the upper surface; basal leaves, including petiole and spines 30–33 × 6–7 cm (petiole 10 cm long), lamina lanceolate-oblong in outline, attenuate towards the base, sinuate-lobate, lobes triangular, alternately large and small, with 4–7 mm long, rigid spines at the margin, midrib thick; lower stem leaves similar to the basal leaves but smaller; middle stem leaves sessile, cordate-auriculate at the base, not decurrent, sub-amplexicaul; upper leaves gradually reduced, shallowly lobed, spinulose along the margin, the uppermost much diminished, bract-like. Capitula solitary, more or less 20-flowered; peduncle 4–10 mm long. Involucre densely arachnoid, oblong, after flowering time campanulate, rounded and rarely cuneate at base, c. 15 long, 7–14 mm wide; phyllaries ca. 60, 7–8 seriate, erect, regularly imbricate, all similar, acicular; outers and median bracts shorter than the inner phyllaries; innermost linear-oblong, acuminate, slightly exceeding, outside papillose, sometimes purple; receptacle bristles smooth; corolla pink, c. 15 mm long, limb 8 mm, tube 7 mm and lobes 2 mm long; anther tube pink, glabrous. Mature achenes light brown, obovoid, attenuate toward the base, truncate above, almost compressed, longitudinally striate, indistinctly scabrous, dark maculate, 5–6 × 3–4 mm; pappus caducous (Fig. 3).

**Etymology.** — The specific epithet refers to the German botanist, Joseph Friedrich Nicolaus Bornmüller (1862-1948).

**Distribution.** — *Cousinia bornmulleriana* is known from a small area in the Avallan Mountains of Kamyaran road to Sanandaj in Kordestan province. It was found at altitudes above 1950 m (Fig. 4).

**Habitat and ecology.** — The area of the type specimen is about 1 km<sup>2</sup> and mountainous. The species that are associated with this species in the area are including *Scutellaria condensata* Rech.f., *Parietaria lusitanica* L., *Phyllocara aucheri* (DC.) Gusuleac, and *Cerasus* sp.

#### Diagnostic key of the new species and its relatives

1. Capitula sessile; synflorescence spike-like ..... *C. haussknechtii* C.Winkl.
- Capitula peduncled; synflorescence a large corymbose ..... 2
2. Phyllaries ±60, erect-imbricate; flowers ±20 ..... *C. bornmulleriana* Attar & Maroofi
- Phyllaries ±160, erect-imbricate; flowers ±60 ..... *C. karkasensis* Mehregan & Djavadi

#### Examined specimens

*C. haussknechtii* C.Winkl.: **Iran.** Kordestan [Kordestan Province]: Shahu and Ovrman Mt., 2100-2400 m, *Haussknecht* 569 (Holotype, G,

image! G00223447); Kordestan [Kordestan Province], mont. Kuh-i-Gerru, 2.8.1909, *Strauss* (JE, image! JE00034787); [Kordestan Province] Garrin (Gerru) Mount., 2.8.1908, *Strauss* (JE, image! JE00034788); [Kordestan Province] Garrin (Gerru) Mount., 2.8.1908, *Strauss* (JE, image! JE00034789); [Kordestan Province]: 40 km from Marivan to Paveh, Zhalaneh passage, 1800-2300 m, *Attar, Maroufi, Hamzeh* 46002 (TUH).

*C. karkasensis* Mehregan & Djavadi: **Iran.** [Esfahan Province]: Natanz, Tameh village, Mt. Karkas, 2300 m, 10.08.2003, *Mehregan* 96022 (TARI, image!).


#### *Cousinia semnanensis*

Attar, Mirtadzadini & Sotoodeh, **sp. nov.** (Fig. 5)


**Diagnosis.** — *Cousinia semnanensis* belongs to sect. *Stenocephalae* and related to *C. lucida*, *C. cylindracea*, *C. glaucopsis*, and *C. alexeenkoana* (as shown in the diagnostic key and table 2), but differs from them based on the characters such as habit, capitulum shape, flower and bract numbers, and leaves' position on the stem.

**TYPE:** Iran. Semnan Province: 12 km after Shahmirzad to Chashm, near the stream, 31.7.2014, 2471 m, *Attar, Mirtadzadini, Rastegar* 48707 (Holotype, TUH; Isotype, TARI).

Perennial, 40–50 cm height, branched from the base. Stems grayish, younger branches arachnoid, then glabrescent, with the remnants of old petioles at the base, interrupted winged, ending in 2–3 capitula. Basal leaves up to 10 cm long (petiole ca. 5 mm long) and 5 cm broad (with spines), elliptic, pinnatifid-pinnatisect, grayish, densely arachnoid on both surfaces; midrib prominent; lobes ending to 3–5 mm long spines; stem leaves sessile, amplexicaul, elliptic, unequally cordate at base, lobed-pinnatifid, terminal lobe long spiny, gradually reduced in the upper part of the stem; uppermost ones close the capitula. Synflorescence racemose-corymbose; capitulum 2 or 3 in each branch, ca. 20-flowered; involucre 15–25 mm in diam., ca. 15 mm long, campanulate, loosely arachnoid; phyllaries 35–40; outers leaf-like, irregular at the margin, equal or longer than capitulum, 20 × 4 mm, lanceolate, long acuminate; medians narrower, lanceolate-subulate, ca. 20 × 2.5 mm; inners white-membranous, oblong-apiculate, 20 × 2 mm; innermost linear, white-membranous; corolla ca. 12 mm long, pale pink; tube ca. 4 mm long; limb ca. 8 mm; anther tube glabrous, concolored. Mature achene 5–6 × 2.5 mm long, obovate, compressed, attenuate toward the base, brown.



**Figure 3.** Floral parts of *Cousinia bornmulleriana* Attar & Maroofi. **a.** habit; **b.** a part of branch; **c.** capitula; **d.** achenes; **e.** corolla; **f.** bristles.



**Figure 4.** Distribution map of *Cousinia bornmulleriana* Attar & Maroofi, sp. nov. (red), *C. karkasensis* (black) and *C. haussknechtii* (blue) in Iran.

**Paratype:** Semnan Province: Chashm road, Hey-Kuh, 3108 m, 31.7.2014, Attar, Mirtadzadini & Rastegar s.n. (Paratype, IRAN).

**Etymology.** — The specific epithet refers to the Semnan province in North-East Iran.

**Distribution.** — This endemic species is distributed as a solitary individual in the Hey-Kuh region of

Semnan province at an altitude of ca. 2500 meters above sea level (Fig. 6).

**Habitat and ecology.** — This species is distributed in mountainous regions at altitudes ranging from 2400 to 3000 m, and sometimes beside rivers.

**Table 1.** Morphological comparison of *Cousinia bornmulleriana* and *C. haussknechtii*.


	<i>Cousinia bornmulleriana</i>	<i>Cousinia Haussknechtii</i>
Inflorescence	large panicle	Racemose
Peduncle length	thin, 4–10 mm	thick, 1–3 mm, subsessile
Capitula, capitula base	7–14 mm, connate	20–25 mm, rounded
Phyllaries no.	±60	±100
Flowers no.	±20	±30
Corolla length	±15 mm, pink or purple	12–15 mm, purple
Limb and tube length	7 mm, 8 mm	9 mm, 6 mm
Corolla lobes	2 mm	4–6 mm
Matured achene	5–6 x 3–4 mm, light brown, almost compressed, obovoid, dark-maculate, longitudinally striate, indistinctly scabrous	5.5 x 3 mm, gray-brown, compressed, gray-brown, irregular dark-maculate, indistinctly longitudinally striate

**Figure 5.** Left: *Cousinia semnanensis* Attar, Mirtadzadini & Sotoodeh, sp. nov. (Holotype, Attar, Mirtadzadini, *Rastegar* 48707); middle and Right: capitula and habit.**Diagnostic key of new species and its relatives**

1. Capitula narrow cylindrical, 3–10-flowered, 2–4.5 mm in diam ..... 2  
- Capitula broad campanulate, ca. 20-flowered, 15–25 mm in diam .....  
.....*C. semnanensis* Attar, Mirtadz. & Sotoodeh
2. Capitula up to 10-flowered ..... 3  
- Capitula 10–20-flowered ..... 4
3. Uppermost leaves surrounding heads, usually exceeding capitula, with a long spine at the apex ..... *C. lucida* DC.  
-Uppermost leaves remote the capitula, not as above ..... *C. cylindracea* Boiss.
4. Phyllaries all subequal; flowers 12–14 ..... *C. alexeenkoana* C.Winkl. & Bornm.  
-Phyllaries unequal; flowers 12–16 ..... *C. glaucopsis* Bornm. & Rech.f.

**Examined specimens:**

***Cousinia glaucopsis*** Bornm. & Rech.f.: **Iran.** Tehran [Tehran Province]: Firouz-Kuh, 17 km Firouz-Kuh to Semnan, 2050 m, *Djavadi, Ghaffari & Bakhsheshi* 9015 (IRAN); 20–35 km Firouz-Kuh to Semnan, 1800–2200 m, *Mousavi & Karavar* 9017 (IRAN, 33866-E); 31 km Firouz-Kuh to Semnan, Bashm pass, 2600 m, *Assadi & Mozaffarain* 40330 (TARI); 25 km Firouz-Kuh to Semnan, Gur-Sefid, 2200 m, *Eskandari & Amini Rad* 43479 (IRAN); Firouz-Kuh, Saran-Za village, 2030 m, *Attar, Zare & Saber* 34400 (TUH); Semnan: 13 km Chashm, Chashm pass, 2173 m, 34398 (TUH); Chashm pass, 2300 m, *Mozaffarian* 78038 (TARI); 35 km Shahmirzad to Foulad Mahalleh, 2300 m, *Assadi & Mozaffarian* 40442 (TARI); Semnan, After to Emamzadeh Abdollah, Eij, 2200–2700 m, *Mozaffarian* 72685 (TARI).



**Figure 6.** Distribution map of *Cousinia semnanensis* Attar, Mirtadzadini & Sotoodeh (red), *C. alexeenkoana* (yellow), *C. glaucopsis* (blue), *C. cylindrica* (green), and *C. lucida* (black).

**Table 2.** Morphological comparison of *Cousinia semnanensis*, *C. alexeenkoana*, *C. glaucopsis*, *C. cylindrica*, and *C. lucida*.

	<i>C. semnanensis</i> Attar, Mirtadzadini & Sotoodeh	<i>C. alexeenkoana</i> Borrm.	<i>C. glaucopsis</i> Borrm. & Rech.f.	<i>C. cylindrica</i> Boiss.	<i>C. lucida</i> DC.
Plant height (cm)	ca. 50	ca. 35	25–35	15–40	15–50
Leaves color and indumentum	grayish, densely araneose, concolous	green, glabrous	green-bluish, araneose-tomentose on both sides, with yellow glands, glabrescent	yellowish-green, glabrous	green on both sides, covered with glandular hairs, punctate
Uppermost leaves	closed and surrounding capitula	closed and surrounding capitula	remote the capitula	remote the capitula	closed and surrounding capitula
Involucre size (mm)	15–25 x 15	14–16 (-20) x 5–6 (-7)	14–16 x 5–6	11–13 x 3–5	12–13 x 2.5–3
Involucre shape	broad campanulate	broad cylindric, narrower at the base	cylindric, narrower at the base	narrow cylindric	narrow cylindric
Phyllaries no.	35–40	16–35	35–40	22–35	22–30
Phyllaries size	all subequal or outer slightly longer	all subequal	outers shorter than the others	outers shorter than the others	outers shorter than the others, recurved at tip
Phyllaries outside and margin	leaf-like, irregular at the margin	ciliate-serrulate	ciliate along the margin, outside in upper half glandular-papillose	ciliate along the margin, outside especially in upper half glandular-papillose,	covering with glandular and papillae hairs in outside and along the margin, ciliate along the margin
Innermost phyllaries	Linear	under apex ciliated, cuspidate	widened toward apex, acuminate	apiculate, cartilaginous along the margin	widened toward apex, acuminate, cartilaginous along the margin
Corolla color	pale pink	yellow	golden yellow	yellow	yellow
Corolla no.	ca. 20	12–14	11–16	5–7	4–5
Corolla, limb, and tube length (mm)	12, limb 8, tube 4	10–13, limb 8, two times longer than the tube, tube 4	11–13.5, limb 7.5–8.5, tube 3.5–5	9.5–10.5, limb 7.5, tube 2–3	ca. 9, limb ca. 7, tube 1.5–2
Achene (mm)	5–6 x 2.5	5 x 3	4 x 1.5	4 x 1.8	2–2.5 x 1–1.5
Achene ornamentation	obovate, compressed, attenuate toward the base, brown	5-costate	truncate-shortly dentate above, brown	linear, costate, almost dentate above, brown-maculate	costate, rugose, distinctly dentate above, dark brown

***Cousinia cylindracea* Boiss.: Iran.** [Markazi Province]: Saveh, Nobaran to Tafresh, between Kahlou and Fark, 1800 m, *Mozaffarian* 72081 (TARI); Arak, Anjadan, 2000-2400 m, *Mozaffarian* 63808 (TARI); Arak, Khan-e Miran, Sefid Khani mountain, 2100-2600 m, *Mozaffarian* 48216 (TARI); Arak, Toureh, Besri, NE slopes of Alandagh mountain, 2100-3100 m, *Mozaffarian* s.n. (TARI); 6 km S Arak, Senejan, Farajollah, *Akhani* 242 (HSBU); Arak, Latedar, 2100-2600 m, *Assadi* 75065 (TARI); Arak, Shahzand, Hafteh-o Emarat, Alim-abad, Alvand mountain, 2100-2800 m, *Mozaffarian* 63892 (TARI); Arak, Moudar, *Djavadi* 53183 (IRAN); Hamedan [Hamedan Province]: Boroujerd, Ghahreman & Sheykhoh-Eslam 11660 (TUH); Boroujerd, Garrin mountain, 2060-2500 m, *Assadi* & *Mehregan* 89090 (TARI); 40 km Boroujerd via Arak road, Safar-Ali village, 2200 m, *Mozaffarian* & *Sardabi* 42169 (TARI); Alvand, 1500 m, *Mirdamadi* 001376 (UTFH); Silvar, *Gauba* & *Sabeti* 111379 (UTFH); Kaboutar Ahang, Shirinsou, 1400 m, *Djavadi* & *Ghanbari* 20379 (IRAN); Asad-abad pass, Gharavol Khani mountain, 2500-2600 m, *Mozaffarian* 64982 (TARI); *Asad-abad*, *Sabeti* 8919 (IRAN, 34014-E); Asad-abad, 1780 m, *Djavadi* & *Ghanbari* 20378 (IRAN); Asad-abad, *Abaei* & *Mojib* 8921 (IRAN); Asad-abad to Gharavol Khaneh mountain, 2500-2700 m, *Mozaffarian* 6498 (TARI); Hamedan, Shahnaz dam, Yalfan, 1980-2200 m, 8756 (IRAN, 33888-E); Lorestan [Lorestan Province]: Khorram-abad, Sefid Dasht to Shoulabad, 2200 m, *Attar* & *Mirtaj* 24907 (TUH); Bisheh, 1948, *Rechinger* & *Manouchehri* 8925 (IRAN, 5745-E); Kuh-e Rig, *Rechinger* & *Mousavi* 8925 (IRAN); Kuh-e Rig, opposite to Kohian, 2000-24000 m, *Mozaffarian* s.n. (TARI); Alastar, Dartang, Garrin mountain, 2130-2800 m, *Assadi* & *Mehregan* 88907 (TARI); 5 km W Azna, 1930 m, *Pabot* 2078 (TARI); Oshtoran Kuh, Tihoun village, 2000-2500 m, *Assadi* & *Mozaffarian* 37052 (TARI); Kordestan [Kordestan Province]: Kamiaran, Bavaneh village, 1720-1900 m, *Assadi* 78814 (TARI); 17 km Sanandaj to Kamiaran, 1500 m, *Assadi* s.n. (TARI); Sanandaj, Chehel Cheshmeh, Saral, 2000-2350 m, *Mozaffarian* s.n. (TARI); Saral, *Mozaffarian* 71573 (TARI); 25 km Sanandaj to Dehgolan, 2040 m, *Mozaffarian* 83778 (TARI); Sanandaj to Qorveh, Salavat-abad pass, 2000-2400 m, *Assadi* s.n. (TARI); 22 km Sanandaj to Marivan, Tijtj, 2070 m, *Mozaffarian* 87373 (TARI); Esfahan: Samirom, 2500 m, *Ghahreman*, *Attar* & *Mehdigholi* 25116 (TUH); Semirom, 12 km Hanna to Komme, 2350 m, *Mozaffarian* & *Maassoumi* s.n. (TARI); Kashan, Qamsar toward Ghahroud road to Esfahan, after Reza-abad, 3100 m, *Assadi*, *Jamzad* & *Azizian* 80047 (TARI); Kashan, between Qamsar and Reza-

abad, 2450 m, *Assadi*, *Jamzad* & *Azizian* 80053 (TARI).

***Cousinia lucida* DC.: Iran.** Lorestan [Lorestan Province]: Doroud to Azna, 25 km Azna, Ghahreman & Attar 21850 (TUH); Doroud, Saravand village, 2100 m, *Mozaffarian* & *Sardabi* 42398 (TARI); Aligoudarz to Shoul-abad, 2300 m, *Runemark* & *Lazari* 26521 (TARI); Hamadan [Hamedan Province]: Alvand Mountain, Babaei 9080 (IRAN); Hamadan to Toyserkan, 12 km Ganj Nameh, 2400 m, *Termeh*, *Mousavi* & *Tehrani* 9083 (IRAN); Esfahan [Esfahan Province]: Ghamishlou protected region, Hazarmani, 2100 m, *Yousefi* 1702 (TARI).

***Cousinia alexeenkoana* Bornm.: Iran.** Qazvin [Qazvin Province]: Qazvin to Hamedan, Avaj, 2100 m, *Assadi* & *Mozaffarian* 36697 (TARI); 48 km Qazvin to Zanjan, *Foroughian* & *Hariri* 4173 (TARI); Hamedan: after Kaboutar-Ahang, Ghondouz, 1860 m, *Ghahreman* & *Attar* TUH-20555; Kaboutar-Ahang toward Shirinsou, 1400 m, 27-30.6.1998, *Djavadi* & *Ghanbari* IRAN-20377; Hamedan, Soltan-Boulagh, between Avaj and Razan, Kharaghan mountains, 1750 m, 27-30.6.1998, *Djavadi* & *Ghanbari* IRAN-20376; Razan to Avaj, 7-8 km to Avaj, 2000-2100 m, *Termeh*, *Eskandari* & *Falsavi* IRAN-43527; Zanjan [Zanjan Province]: Zanjan to Hamedan, Takht Mountains, 2150-2250 m, 5.7.1974, *Termeh* & *Mousavi* IRAN-8757 (33894-E); Esfahan [Esfahan Province]: Natanz, Karkas mountain, 2700 m, *Foroughi* 4047 (TARI); Karkas mountain, 2600 m, *Mozaffarian* & *Maassoumi* 78351 (TARI).

## ACKNOWLEDGEMENTS

The authors would like to thank the reviewers for their time and their valuable comments. We also thank Dr. John Edmonson (Kew, UK) for his time, corrections and useful comments.

## REFERENCES

- Assadi, M.** 2009. Four new species of the genus *Cousinia* Cass. (Asteraceae) from Iran. *Iranian Journal of Botany* 15: 36-44.
- Assadi, M.** 2010. A taxonomic revision of *Cousinia* Cass. sect. *Chaetocephalae* (Asteraceae). *Iranian Journal of Botany* 16: 191-196.
- Assadi, M.** 2011. Two new species of the genus *Cousinia* (Asteraceae) from Iran. *Iranian Journal of Botany* 17: 6-9.
- Atazadeh, N., Sheidai, M., Attar, F., Ghahremaninejad, F. & Koohdar, F.** 2020. A palynological study of genus *Cousinia* Cass. (Family Asteraceae), sections

- Cynaroideae* Bunge and *Platyacanthae* Rech. f. Grana 59: 428-443. DOI: 10.1080/00173134.2020.1781247
- Attar, F.** 2011. New taxa of *Cousinia* Cass., sect. *Stenocephalae* Bunge (Asteraceae) from Iran. Journal of Taxonomy and Biosystematics 3: 59-68.
- Attar, F. & Amini Rad, M.** 2019. *Cousinia sharifii*, a new species of *Cousinia* sect. *Cynaroideae* (Asteraceae) from NW Iran. Phytotaxa 395: 287-295.
- Attar, F. & Djavadi, S.B.** 2010. A taxonomic revision of *Cousinia*, sect. *Cynaroides* (Asteraceae, Cardueae) in the flora of Iran. Iranian Journal of Botany 16: 130-184.
- Attar, F., Ghahreman, A., Mahdigholi, K. & Sheidai, M.** 2007. A synopsis of *Cousinia* sect. *Serratuloides* (Asteraceae) in Iran. Novon 17: 1-7.
- Attar, F. & Mirtadzadini, M.** 2009. Two new species of *Cousinia* Cass. sect. *Pugioniferae* Bunge (Asteraceae, Cardueae), from east and southeast of Iran. Iranian Journal of Botany 15: 146-152.
- Attar, F., Mozaffarian, V. & Mirtadzadini, M.** 2016. *Cousinia brevicaulis* sp. nov. (sect. *Lasiandra*) for the flora of Iran. Nordic Journal of Botany 34: 174-177.
- Djavadi, S.B. & Attar, F.** 2006. Two new species of *Cousinia* Cass. of sect. *Stenocephalae* Bunge from Iran. Feddes Repertorium 117: 453-458.
- Huber-Morath, A.** 1975. *Cousinia*. In: Davis, P.H., (editor), Flora of Turkey and the East Aegean Islands. vol. 5: 329-353. Edinburgh University Press, Edinburgh.
- López-Vinyallonga, S., Mehregan, I., Garcia-Jacas, N., Tscherneva, O., Susanna, A. & Kadereit, J.W.** 2009. Phylogeny and evolution of the *Arctium-Cousinia* complex (Compositae, Cardueae-Carduinae). Taxon 58: 153-171. DOI: 10.1002/tax.581016
- Mehregan, I. & Assadi, M.** 2010. *Cousinia* sect. *Argenteae* (Asteraceae, Cardueae), a new section including a new species from NE Iran. Willdenowia 39: 265-271.
- Mehregan, I., Djavadi, S.B. & Pahlevani, A.H.** 2010. *Cousinia* (sect. *Hausknechtianae*) *karkasensis*, a new species from Karkas Mts. in central Iran. Iranian Journal of Botany 16: 200-203.
- Mirtadzadini, M. & Attar, F.** 2014. *Cousinia kermanensis* and *C. pulcherantha* spp. nov. (Compositae, sect. *Congestae* and sect. *Leiocaulis*) from southeastern Iran. Nordic Journal of Botany 32: 279-284.
- Rastegar, A., Attar, F. & Mirtadzadini, M.** 2018. Novelities in Iranian *Cousinia* sect. *Cynaroideae* (Asteraceae, Cardueae): new taxa and taxonomic notes. Phytotaxa 343: 227. DOI: 10.11646/phytotaxa.343.3.3
- Rechinger, K.H.** 1972. Compositae-Cynareae I: *Cousinia*. In: Rechinger, K.H. (editor), Flora Iranica, vol. 90: 1-329. Akademische Druck- und Verlagsanstalt, Graz.
- Rechinger, K.H.** 1979. Compositae-Cynareae III. *Cousinia*. In: Rechinger, K.H. (editor), Flora Iranica, vol. 139: 108-153. Akademische Druck- und Verlagsanstalt, Graz.
- Rechinger, K.H.** 1986. *Cousinia*: morphology, taxonomy, distribution and phytogeographical implications. Proceedings of the Royal Society of Edinburgh. Section B. Biological Sciences 89: 45-58. DOI: 10.1017/S0269727000008897
- Sennikov, A.N.** 2013. A revision of *Cousinia* sections *Alpinae* (syn. *Carduncellus*), *Subappendiculatae* and *Tianschanicae* (Asteraceae) in the Kirghizian Tian-Shan and the neighbouring territories. Phytotaxa 5: 1. DOI: 10.11646/phytotaxa.5.1.1
- Tscherneva, O.V., Joharchi, M. & Ghahremaninejad, F.** 2005. A new species of the genus *Cousinia* (Asteraceae) from Iran. Botanicheskii Zhurnal (Moscow & Leningrad [St. Petersburg]) 90: 411-414.
- Tscherneva, O.V.** 1962. *Cousinia*. In: Shishkin, B.K. (editor), Flora of the U.S.S.R., vol. 27. Academy of Sciences of the U.S.S.R., Leningrad, pp. 108-357.

\*\*\*\*\*

**How to cite this article:**

**Attar, F., Maroofi, H., Mirtadzadini, M. & Sotoodeh, A.** 2022. Two new species of genus *Cousinia* (Asteraceae, Cardueae) from W and NE Iran. Nova Biologica Reperta 9: 144-152.